

A Package to Integrate Rational Functions using the Minimal Algebraic Extension to the Constant Field

Neil Langmead
Konrad-Zuse-Zentrum für Informationstechnik (ZIB)
Takustrasse 7
D- 14195 Berlin Dahlem
Berlin Germany

January 1997

Package to Integrate Rational Functions using the minimal algebraic extension to the constant field

Contents

1	Rational Integration	3
1.1	Syntax of <i>ratint</i>	3
1.2	Examples	3
2	The Algorithm	5
3	The log_sum operator	7
3.1	More examples	8
4	Options	8
4.1	LogtoAtan function	9
5	Hermite's method	10
6	Tracing the <i>ratint</i> program	10
7	Bugs, suggestions and comments	11

1 Rational Integration

This package implements the Horowitz/ Rothstein/ Trager algorithms [Ged92] for the integration of rational functions in REDUCE. We work within a field K of characteristic 0 and functions $p, q \in K[x]$. K is normally the field Q of rational numbers, but not always. These procedures return $\int \frac{p}{q} dx$. The aim is to be able to integrate any function of the form p/q in x , where p and q are polynomials in the field Q . The algorithms used avoid algebraic number extensions wherever possible, and in general, express the integral using the minimal algebraic extension field.

1.1 Syntax of *ratint*

This function has the following syntax:

ratint(p,q,var)

where p/q is a rational function in var . The output of *ratint* is a list of two elements: the first is the polynomial part of the integral, the second is the logarithmic part. The integral is the sum of these parts.

1.2 Examples

consider the following examples in REDUCE:

```
ratint(1,x^2-2,x);
```

$$\left\{ 0, \frac{\log\left(\frac{\sqrt{2}x-2}{\sqrt{2}}\right) - \log\left(\frac{\sqrt{2}x+2}{\sqrt{2}}\right)}{2\sqrt{2}} \right\}$$

```
p:=441*x^7+780*x^6-2861*x^5+4085*x^4+7695*x^3+3713*x^2-43253*x
+24500;
```

```
q:=9*x^6+6*x^5-65*x^4+20*x^3+135*x^2-154*x+49;
```

```
ratint(p,q,x);
```

$$\left\{ \frac{49}{2}x^6 + \frac{6}{147}x^5 - \frac{226}{49}x^4 + \frac{268}{49}x^3 + \frac{1332}{49}x^2 - \frac{2809}{147}x + \frac{752}{21} - \frac{256}{9}, 0 \right\}$$

```
k:=36*x^6+126*x^5+183*x^4+(13807/6)*x^3-407*x^2-(3242/5)*x+(3044/15);
l:=(x^2+(7/6)*x+(1/3))^2*(x-(2/5))^3;
```

```
ratint(k,l,x);
```

$$\frac{5271}{5} \left(x^3 + \frac{39547}{52710} x^2 - \frac{31018}{26355} x + \frac{7142}{26355} \right) \left\{ \frac{x^4}{30} + \frac{11}{25} x^3 - \frac{11}{25} x^2 - \frac{2}{25} x + \frac{4}{75} \right\}$$

$$- \frac{37451}{16} \left(\log(x - \frac{2}{5}) + \frac{91125}{37451} \log(x + \frac{2}{3}) - \frac{128000}{37451} \log(x + \frac{1}{2}) \right)$$

```
ratint(1,x^2+1,x);
```

$$\{0, \log_sum(\beta, \beta^2 + \frac{1}{4}, 0, \log(2\beta x - 1)\beta)\}$$

The meaning of the `log_sum` function will be explained later.

2 The Algorithm

The following main algorithm is used:

```

procedure ratint( $p, q, x$ );
solution_list  $\leftarrow$  HorowitzReduction( $p, q, x$ )
 $c/d \leftarrow$  part(solution_list,1)
poly_part  $\leftarrow$  part(solution_list,2)
rat_part  $\leftarrow$  part(solution_list,3)
rat_part  $\leftarrow$  LogarithmicPartIntegral(rat_part,  $x$ )
return(rat_part +  $c/d$  + poly_part)
end

```

The algorithm contains two subroutines, *HorowitzReduction* and *rt*. *HorowitzReduction* is an implementation of Horowitz' method to reduce a given rational function into a polynomial part and a logarithmic part. The integration of the polynomial part is a trivial task, and is done by the *int* operator in REDUCE. The integration of the logarithmic part is done by the routine *rt*, which is an impementation of the Rothstein and Trager method. These two answers are outputted in a list, the complete answer being the sum of these two parts.

These two algorithms are as follows:

```

procedure how( $p, q, x$ )

```

for a given rational function p/q in x , this algorithm calculates the reduction of $\int(p/q)$ into a polynomial part and logarithmic part.

```

poly_part  $\leftarrow$  quo( $p, q$ );  $p \leftarrow$  rem( $p, q$ );
 $d \leftarrow$  GCD( $q, q'$ );  $b \leftarrow$  quo( $q, d$ );  $m \leftarrow$  deg( $b$ );
 $n \leftarrow$  deg( $d$ );
 $a \leftarrow \sum_{i=1}^{m-1} a_i x^i$ ;  $c \leftarrow \sum_{i=1}^{n-1} c_i x^i$ ;

```

```

 $r \leftarrow b * c' - \text{quo}(b * d', d) + d * a$ ;

```

```

for  $i$  from 0 to  $m + n - 1$  do

```

```

    {
         $eqns(i) \leftarrow$  coeff( $p, i$ ) = coeff( $r, i$ );
    };

```

```

solve(eqns, { $a(0), \dots, a(m-1), c(0), \dots, c(n-1)$ });

```

```

return( $c/d + \int$  poly_part +  $\int a/b$ );

```

```

end;

```

```

procedure RothsteinTrager( $a, b, x$ )
% Given a rational function  $a/b$  in  $x$  with  $\deg(a) < \deg(b)$ , with  $b$  monic and square free, we
calculate  $\int(a/b)$ 
 $R(z) \leftarrow \text{residue}(a - zb', b)$ 
 $(r_1(z) \dots r_k(z)) \leftarrow \text{factors}(R(z))$ 
integral  $\leftarrow 0$ 

for  $i$  from 1 to  $k$  do
  {
     $d \leftarrow \text{degree}(r_i(z))$ 
    if  $d = 1$  then
      {
         $c \leftarrow \text{solve}(r_i(z) = 0, z)$ 
         $v \leftarrow \text{GCD}(a - cb', b)$ 
         $v \leftarrow v/\text{lcoeff}(v)$ 
        integral  $\leftarrow \text{integral} + c * \log(v)$ 
      }
    else {
      % we need to do a GCD over algebraic number field
       $v \leftarrow \text{GCD}(a - \alpha * b', b)$ 
       $v \leftarrow v/\text{lcoeff}(v)$ , where  $\alpha = \text{root\_of}(r_i(z))$ 
      if  $d=2$  then {
        % give answer in terms of radicals
         $c \leftarrow \text{solve}(r_i(z) = 0, z)$ 
        for  $j$  from 1 to 2 do {
           $v[j] \leftarrow \text{substitute}(\alpha = c[j], v)$ 
          integral  $\leftarrow \text{integral} + c[j] * \log(v[j])$ 
        }
      }
      else {
        % Need answer in terms of root_of notation
        for  $j$  from 1 to  $d$  do {
           $v[j] \leftarrow \text{substitute}(\alpha = c[j], v)$ 
          integral  $\leftarrow \text{integral} + c[j] * \log(v[j])$ 
          % where  $c[j] = \text{root\_of}(r_i(z))$  }
        }
      }
    }
  }
return(integral)
end

```

3 The log_sum operator

The algorithms above returns a sum of terms of the form

$$\sum_{\alpha | R(\alpha)=0} \log(S(\alpha, x)),$$

where $R \in K[z]$ is square free, and $S \in K[z, x]$. In the cases where the degree of $R(\alpha)$ is less than two, this is merely a sum of logarithms. For cases where the degree is two or more, I have chosen to adopt this notation as the answer to the original problem of integrating the rational function. For example, consider the integral

$$\int \frac{a}{b} = \int \frac{2x^5 - 19x^4 + 60x^3 - 159 + x^2 + 50x + 11}{x^6 - 13x^5 + 58x^4 - 85x^3 - 66x^2 - 17x + 1} dx$$

Calculating the resultant $R(z) = \text{res}_x(a - zb', b)$ and factorising gives

$$R(z) = -190107645728000(z^3 - z^2 + z + 1)^2$$

Making the result monic, we have

$$R_2(z) = z^3 - z^2 + z + 1$$

which does not split over the constant field Q . Continuing with the Rothstein Trager algorithm, we now calculate

$$\gcd(a - \alpha b', b) = z^2 + (2 * \alpha - 5) * z + \alpha^2,$$

where α is a root of $R_2(z)$.

Thus we can write

$$\int \frac{a}{b} = \sum_{\alpha | \alpha^3 - \alpha^2 + \alpha + 1 = 0} \alpha * \log(x^2 + 2\alpha x - 5x + \alpha^2),$$

and this is the answer now returned by REDUCE, via a function called *log_sum*. This has the following syntax:

$$\text{log_sum}(\alpha, \text{eqn}(\alpha), 0, \text{sum_term}, \text{var})$$

where α satisfies $\text{eqn} = 0$, and *sum_term* is the term of the summation in the variable *var*. Thus in the above example, we have

$$\int \frac{a}{b} dx = \text{log_sum}(\alpha, \alpha^3 - \alpha^2 + \alpha + 1, 0, \alpha * \log(x^2 + 2\alpha x - 5x + \alpha^2), x)$$

Many rational functions that could not be integrated by REDUCE previously can now be integrated with this package. The above is one example; some more are given on the next page.

3.1 More examples

$$\int \frac{1}{x^5 + 1} dx = \frac{1}{5} \log(x + 1) + 5 \log_sum(\beta, \beta^4 + \frac{1}{5}\beta^3 + \frac{1}{25}\beta^2 + \frac{1}{125}\beta + \frac{1}{625}, 0, \log(5 * \beta + x) * \beta)$$

which should be read as

$$\int \frac{1}{x^5 + 1} dx = \frac{1}{5} \log(x + 1) + \sum_{\beta | \beta^4 + \frac{1}{5}\beta^3 + \frac{1}{25}\beta^2 + \frac{1}{125}\beta + \frac{1}{625} = 0} \log(5 * \beta + x) \beta$$

$$\int \frac{7x^{13} + 10x^8 + 4x^7 - 7x^6 - 4x^3 - 4x^2 + 3x + 3}{x^{14} - 2x^8 - 2x^7 - 2x^4 - 4x^3 - x^2 + 2x + 1} dx = \log_sum(\alpha, \alpha^2 - \alpha - \frac{1}{4}, 0, \log(-2\alpha x^2 - 2\alpha x + x^7 + x^2 - 1) * \alpha, x),$$

$$\int \frac{1}{x^3 + x + 1} dx = \log_sum(\beta, \beta^3 - \frac{3}{31}\beta^2 - \frac{1}{31}, 0, \beta \log(-\frac{62}{9}\beta^2 + \frac{31}{9}\beta + x + \frac{4}{9})).$$

4 Options

There are several alternative forms that the answer to the integration problem can take. One output is the *log_sum* form shown in the examples above. There is an option with this package to convert this to a "normal" sum of logarithms in the case when the degree of *eqn* in α is two, and α can be expressed in surds. To do this, use the function *convert*, which has the following syntax:

convert(exp)

If *exp* is free of *log_sum* terms, then *exp* itself is returned. If *exp* contains *log_sum* terms, then α is represented as surds, and substituted into the *log_sum* expression. For example, using the last example, we have in REDUCE:

```
2: ratint(a,b,x);
```

```
{0,
```

```
      2      1
log_sum(alpha,alpha  - alpha - ---,0,
```

```
      2      7      2
log( - 2*alpha*x  - 2*alpha*x + x  + x  - 1)*alpha,x)}
```

```
3: convert(ws);
```


$$\begin{aligned}
& \frac{1}{2} * (\sqrt{2} * \log(-\sqrt{2} * x^2 - \sqrt{2} * x + x^7 - x - 1) \\
& - \sqrt{2} * \log(\sqrt{2} * x^2 + \sqrt{2} * x + x^7 - x - 1) \\
& + \log(-\sqrt{2} * x^2 - \sqrt{2} * x + x^7 - x - 1) \\
& + \log(\sqrt{2} * x^2 + \sqrt{2} * x + x^7 - x - 1))
\end{aligned}$$

4.1 LogtoAtan function

The user could then combine these to form a more elegant answer, using the switch combinelogs if one so wished. Another option is to convert complex logarithms to real arctangents [Bron97], which is recommended if definite integration is the goal. This is implemented in REDUCE via a function *convert_log*, which has the following syntax:

convert_log(exp),

where *exp* is any log_sum expression.

The procedure to convert complex logarithms to real arctangents is based on an algorithm by Rio-boo. Here is what it does:

Given a field K of characteristic 0 such that $\sqrt{-1} \notin K$ and $A, B \in K[x]$ with $B \neq 0$, return a sum f of arctangents of polynomials in $K[x]$ such that

$$\frac{df}{dx} = \frac{d}{dx} i \log\left(\frac{A + iB}{A - iB}\right)$$

Example:

$$\int \frac{x^4 - 3 * x^2 + 6}{x^6 - 5 * x^4 + 5 * x^2 + 4} dx = \sum_{\alpha | 4\alpha + 1 = 0} \alpha \log(x^3 + 2\alpha x^2 - 3x - 4\alpha)$$

Substituting $\alpha = i/2$ and $\alpha = -i/2$ gives the result

$$\frac{i}{2} \log\left(\frac{(x^3 - 3x) + i(x^2 - 2)}{(x^3 - 3x) - i(x^2 - 2)}\right)$$

Applying logtoAtan now with $A = x^3 - 3x$, and $B = x^2 - 2$ we obtain

$$\int \frac{x^4 - 3 * x^2 + 6}{x^6 - 5 * x^4 + 5 * x^2 + 4} dx = \arctan\left(\frac{x^5 - 3x^3 + x}{2}\right) + \arctan(x^3) + \arctan(x),$$

and this is the formula which should be used for definite integration.

Another example in REDUCE is given below:

```

1: ratint(1,x^2+1,x);

*** Domain mode rational changed to arnum

          2    1
{0,log_sum(beta,beta + ---,0,log(2*beta*x - 1)*beta)}
          4

13: part(ws,2);

          2    1
log_sum(beta,beta + ---,0,log(2*beta*x - 1)*beta)
          4

14: on combinelogs;

15: convertlog(ws);

  1      - i*x + 1
---*log(-----)*i
  2      i*x + 1

logtoAtan(-x,1,x);

2*atan(x)

```

5 Hermite's method

The package also implements Hermite's method to reduce the integral into its polynomial and logarithmic parts, but occasionally, REDUCE returns the incorrect answer when this algorithm is used. This is due to the REDUCE operator `pf`, which performs a complete partial fraction expansion when given a rational function as input. Work is presently being done to give the `pf` operator a facility which tells it that the input is already factored. This would then enable REDUCE to perform a partial fraction decomposition with respect to a square free denominator, which may not necessarily be fully factored over \mathbb{Q} .

For a complete explanation of this and the other algorithms used in this package, including the theoretical justification and proofs, please consult [Ged92].

6 Tracing the *ratint* program

The package includes a facility to trace in some detail the inner workings of the *ratint* program. Messages are given at the key stages of the algorithm, together with the results obtained. These messages are displayed when the switch *tracerratint* is on, which is done in REDUCE with the command

```
on traceratint;
```

This switch is off by default. Here is an example of the output obtained with this switch on:

```
Loading image file: /silo/tony/red/lisp/psl/solaris/red/reduce.img
REDUCE Development Version, 21-May-97 ...
```

```
1: load_package ratint;
```

```
2: on traceratint;
```

```
3: ratint(1+x,x^2-2*x+1,x);
```

```
performing Howoritz reduction on  $\frac{x + 1}{x^2 - 2x + 1}$ 
```

```
Howoritz gives:  $\left\{\frac{-2}{x-1}, 0, \frac{1}{x-1}\right\}$ 
```

```
computing Rothstein Trager on  $\frac{1}{x-1}$ 
```

```
integral in Rothstein T is  $\log(x - 1)$ 
```

```
 $\left\{\frac{-2}{x-1}, \log(x - 1)\right\}$ 
```

7 Bugs, suggestions and comments

This package was written when the author was working as a placement student at ZIB Berlin. All comments should therefore be reported to Winfried Neun, ZIB, Takustrasse 7, D 14195 Berlin Dahlem, Germany (email: neun@zib.de).

References

- [Bron97] Bronstein, Manuel, *Symbolic Integration I: Transcendental Functions*, Springer-Verlag, Heidelberg, 1997.
- [Dav88] Davenport, James H. et al, *Computer Algebra- Systems and Algorithms for Algebraic Computation*, Academic Press, 1988.
- [Ged92] Geddes, K.O. et al, *Algorithms for Computer Algebra*, Kluwer Academic Publishers, 1992.
- [Red36] Hearn, Anthony C. and Fitch, John F. *REDUCE User's Manual 3.6*, RAND Corporation, 1995